

Primary Prostate Cryoablation

1. Babaian RJ, Donnelly B and Thrasher JB, et. al. Best practice statement on cryosurgery for the treatment of localized prostate cancer. *J Urol.* 2009; 180: 1993-2004.
2. Malcolm JB, Fabrizio MD and Schellhammer PF, et. al. Quality of life after open or robotic prostatectomy, cryoablation or brachytherapy for localized prostate cancer. *J Urol* 2010; 183: 1822-1829.
3. Kimura M, Mouraviev V and Polascik TJ, et. al. Analysis of urinary function using validated instruments and uroflowmetry after primary and salvage prostate cryoablation. *Urology* 2010; 76:1258-65.
4. Cheetham P. and Katz A., et. al. Long-Term Cancer-Specific and Overall Survival for Men Followed More Than 10 Years After Primary and Salvage Cryoablation of the Prostate. *J. of Endourology* 2010; 24(7): 1123–1129.
5. Dhar N., Ward J.F., and Jones J.S. et. al. Primary Full-Gland Prostate Cryoablation in Older Men (> Age of 75 Years): Results from 860 Patients Tracked with the COLD Registry. *BJU International* 2011; 108: 508-512.
6. Cohen J.K., Miller Jr. R.J., and Baust J. et. al. Ten-Year Biochemical Disease Control for Patients with Prostate Cancer Treated with Cryosurgery as Primary Therapy. *Urology* 2008; 71: 515–518.